



Shrinkage Estimators In Inverse Gaussian Regression Model: Subject Review

Farah A. Yonis  Rafal A. Othman 

Department of Informatics & Statistic, College of Computer & Mathematical Science, University of Mosul, Mosul, Iraq

Article information

Article history:

Received November 29, 2021

Revised: April 25, 2022

Accepted: April 28, 2022

Available online June 1, 2022

Keywords: Multicollinearity; biased estimator; inverse Gaussian regression model; Monte Carlo simulation

Correspondence:

Farah.A. Yonis

farah-abd-ul-ghani@uomosul.edu.iq

Abstract

The presence of the high correlation among predictors in regression modeling has undesirable effects on the regression estimating. There are several available biased methods to overcome this issue. The inverse Gaussian regression model (IGRM) is a special model from the generalized linear models. The IGRM is a well-known model in research application when the response variable under the study is skewed data. Numerous biased estimators for overcoming the multicollinearity in IGRM have been proposed in the literature using different theories. An overview of recent biased methods for IGRM is provided. A comparison among these biased estimators allows us to gain an insight into their performance.

DOI: [10.3389/IQJOSS.2022.174331](https://doi.org/10.3389/IQJOSS.2022.174331) ©Authors, 2022, College of Computer Science and Mathematics, University of Mosul.

This is an open access article under the CC BY 4.0 license (<http://creativecommons.org/licenses/by/4.0/>).

1. Introduction

The inverse Gaussian regression model (IGRM) has been widely used in industrial engineering, life testing, reliability, marketing, and social sciences [1-7]. "Specifically, IGRM is used when the response variable under the study is positively skewed [8-10]. When the response variable is extremely skewness, the IGRM is preferable than gamma regression model [11]. In dealing with the IGRM, it is assumed that there is no correlation among the explanatory variables [12-32]. In practice, however, this assumption often not holds, which leads to the problem of multicollinearity. In the presence of multicollinearity, when estimating the regression coefficients for IGRM using the maximum likelihood (ML) method, the estimated coefficients are usually become unstable with a high variance, and therefore low statistical significance [33]. Numerous remedial methods have been proposed to overcome the problem of multicollinearity [34-38]. The ridge regression method [39] has been consistently demonstrated to be an attractive and alternative to the ML estimation method. Ridge regression is a biased method that shrinks all regression coefficients toward zero to reduce the large variance [40].

This done by adding a positive amount to the diagonal of $\mathbf{X}^T \mathbf{X}$. As a result, the ridge estimator is biased but it guaranties a smaller mean squared error than the ML estimator.

In linear regression, the ridge estimator is defined as

$$\hat{\boldsymbol{\beta}}_{Ridge} = (\mathbf{X}^T \mathbf{X} + k \mathbf{I})^{-1} \mathbf{X}^T \mathbf{y}, \quad (1)$$

where \mathbf{y} is an $n \times 1$ vector of observations of the response variable, $\mathbf{X} = (\mathbf{x}_1, \dots, \mathbf{x}_p)$ is an $n \times p$ known design matrix of explanatory variables, $\boldsymbol{\beta} = (\beta_1, \dots, \beta_p)$ is a $p \times 1$ vector of unknown regression coefficients, \mathbf{I} is the identity matrix

with dimension $p \times p$, and $k \geq 0$ represents the ridge parameter (shrinkage parameter). The ridge parameter, k , controls the shrinkage of β toward zero. The OLS estimator can be considered as a special estimator from Eq. (1) with $k = 0$. For larger value of k , the $\hat{\beta}_{Ridge}$ estimator yields greater shrinkage approaching zero [39, 41].

1. Inverse Gaussian regression model

The inverse Gaussian distribution is a continuous distribution with two positive parameters: location parameter, μ , and scale parameter, τ , denoted as $IG(\mu, \tau)$. Its probability density function is defined as

$$f(y, \mu, \tau) = \frac{1}{\sqrt{2\pi y^3 \tau}} \exp \left[-\frac{1}{2y} \left(\frac{y - \mu}{\mu \sqrt{\tau}} \right)^2 \right], \quad y > 0. \quad (2)$$

The mean and variance of this distribution are, respectively, $E(y) = \mu$ and $\text{var}(y) = \tau \mu^3$.

Inverse Gaussian regression model is considered a member of the generalized linear models (GLM) family, extending the ideas of linear regression to the situation where the response variable is following the inverse Gaussian distribution. Following the GLM methodology, Eq. (1) can re-write in terms of exponential family function as

$$f(y, \mu, \tau) = \frac{1}{\tau} \left\{ -\frac{y}{2\mu^2} + \frac{1}{\mu} \right\} + \left\{ -\frac{1}{2} \ln(2\pi y^3) - \frac{1}{2} \ln(\tau) \right\}, \quad (3)$$

where $C(y, \tau) = -(1/2) \ln(2\pi y^3) - (1/2) \ln(\tau)$ and $\frac{y\theta - a(\theta)}{\phi} = \frac{1}{\tau} \left\{ -\frac{y}{2\mu^2} + \frac{1}{\mu} \right\}$. Here, τ represents the dispersion parameter and $1/\mu^2$ represents the canonical link function.

In GLM, a monotonic and differentiable link function connects the mean of the response variable with the linear predictor $\eta_i = \mathbf{x}_i^T \beta$, where \mathbf{x}_i is the i^{th} row of \mathbf{X} and β is a $(p+1) \times 1$ vector of unknown regression coefficients. Because η_i depends on β and the mean of the response variable is a function of η_i , then $E(y_i) = \mu_i = g^{-1}(\eta_i) = g^{-1}(\mathbf{x}_i^T \beta)$.

Related to the IGR, the $\mu = 1/\sqrt{\mathbf{x}_i^T \beta}$. Another possible link function for the IGRM is log link function, $\mu = \exp(\mathbf{x}_i^T \beta)$.

The model estimation of the IGRM is based on the maximum likelihood method (ML). The log likelihood function of the IGRM under the canonical link function is defined as

$$\ell(\beta) = \sum_{i=1}^n \left\{ \frac{1}{\tau} \left[\frac{y_i \mathbf{x}_i^T \beta}{2} - \sqrt{\mathbf{x}_i^T \beta} \right] - \frac{1}{2\tau y_i} - \frac{\ln \tau}{2} - \ln(2\pi y_i^3) \right\}. \quad (4)$$

The ML estimator is then obtained by computing the first derivative of the Eq. (3) and setting it equal to zero, as

$$\frac{\partial \ell(\beta)}{\partial \beta} = \sum_{i=1}^n \frac{1}{2\tau} \left[y_i - \frac{1}{\sqrt{\mathbf{x}_i^T \beta}} \right] \mathbf{x}_i = 0. \quad (5)$$

Unfortunately, the first derivative cannot be solved analytically because Eq. (4) is nonlinear in β . The iteratively weighted least squares (IWLS) algorithm or Fisher-scoring algorithm can be used to obtain the ML estimators of the IGRM parameters. In each iteration, the parameters are updated by

$$\beta^{(r+1)} = \beta^{(r)} + I^{-1}(\beta^{(r)}) S(\beta^{(r)}), \quad (6)$$

where $S(\boldsymbol{\beta}^{(r)})$ and $I^{-1}(\boldsymbol{\beta}^{(r)})$ are $S(\boldsymbol{\beta}) = \partial \ell(\boldsymbol{\beta}) / \partial \boldsymbol{\beta}$ and $I^{-1}(\boldsymbol{\beta}) = \left(-E \left(\partial^2 \ell(\boldsymbol{\beta}) / \partial \boldsymbol{\beta} \partial \boldsymbol{\beta}^T \right) \right)^{-1}$ evaluated at $\boldsymbol{\beta}^{(r)}$, respectively. The final step of the estimated coefficients is defined as

$$\hat{\boldsymbol{\beta}}_{IGRM} = \mathbf{B}^{-1} \mathbf{X}^T \hat{\mathbf{W}} \hat{\mathbf{m}}, \quad (7)$$

where $\mathbf{B} = (\mathbf{X}^T \hat{\mathbf{W}} \mathbf{X})$, $\hat{\mathbf{W}} = \text{diag}(\hat{\mu}_i^3)$, $\hat{\mathbf{m}}$ is a vector where i^{th} element equals to $\hat{m}_i = (1 / \hat{\mu}_i^2) + ((y_i - \hat{\mu}_i) / \hat{\mu}_i^3)$, and $\hat{\mu} = 1 / \sqrt{\mathbf{x}_i^T \hat{\boldsymbol{\beta}}}$. The covariance matrix of $\hat{\boldsymbol{\beta}}_{IGRM}$ equals

$$\text{cov}(\hat{\boldsymbol{\beta}}_{IGRM}) = \left[-E \left(\frac{\partial^2 \ell(\boldsymbol{\beta})}{\partial \boldsymbol{\beta} \partial \boldsymbol{\beta}^T} \right) \right]^{-1} = \tau \mathbf{B}^{-1}, \quad (8)$$

and the MSE equals

$$\begin{aligned} \text{MSE}(\hat{\boldsymbol{\beta}}_{IGRM}) &= E(\hat{\boldsymbol{\beta}}_{IGRM} - \hat{\boldsymbol{\beta}})^T (\hat{\boldsymbol{\beta}}_{IGRM} - \hat{\boldsymbol{\beta}}) \\ &= \tau \text{tr}[\mathbf{B}^{-1}] \\ &= \tau \sum_{j=1}^p \frac{1}{\lambda_j}, \end{aligned} \quad (9)$$

where λ_j is the eigenvalue of the \mathbf{B} matrix and the dispersion parameter, τ , is estimated by [42]

$$\hat{\tau} = \frac{1}{(n-p)} \sum_{i=1}^n \frac{(y_i - \hat{\mu}_i)^2}{\hat{\mu}_i^3}. \quad (10)$$

2. Ridge estimator

In the presence of multicollinearity, the matrix $\mathbf{X}^T \hat{\mathbf{W}} \mathbf{X}$ becomes ill-conditioned leading to high variance and instability of the ML estimator of the IGRM parameters. As a remedy, Månsson and Shukur [43] proposed the IGR ridge estimator (IGRR) as

$$\begin{aligned} \hat{\boldsymbol{\beta}}_{IGRR} &= (\mathbf{X}^T \hat{\mathbf{W}} \mathbf{X} + k\mathbf{I})^{-1} \mathbf{X}^T \hat{\mathbf{W}} \mathbf{X} \hat{\boldsymbol{\beta}}_{IGRM} \\ &= (\mathbf{X}^T \hat{\mathbf{W}} \mathbf{X} + k\mathbf{I})^{-1} \mathbf{X}^T \hat{\mathbf{W}} \hat{\mathbf{v}}, \end{aligned} \quad (11)$$

where $k \geq 0$. The ML estimator can be considered as a special estimator from Eq. (11) with $k = 0$. Regardless of k value, the MSE of the $\hat{\boldsymbol{\beta}}_{IGRR}$ is smaller than that of $\hat{\boldsymbol{\beta}}_{IGRM}$ because the MSE of $\hat{\boldsymbol{\beta}}_{IGRR}$ is equal to [33]

$$\text{MSE}(\hat{\boldsymbol{\beta}}_{IGRR}) = \tau \sum_{j=1}^p \frac{\lambda_j}{(\lambda_j + k)^2} + k^2 \sum_{j=1}^p \frac{\alpha_j}{(\lambda_j + k)^2}, \quad (12)$$

where α_j is defined as the j^{th} element of $\gamma \hat{\boldsymbol{\beta}}_{IGRM}$ and γ is the eigenvector of the $\mathbf{X}^T \hat{\mathbf{W}} \mathbf{X}$ matrix. Comparing with the MSE of Eq. (9), $\text{MSE}(\hat{\boldsymbol{\beta}}_{IGRR})$ is always small for $k > 0$.

3. Liu estimator

Another popular biased estimator which is known as Liu estimator has been adopted in Poisson regression model. The inverse Gaussian Liu estimator (IGLE) is defined as

$$\hat{\boldsymbol{\beta}}_{IGLE} = (\mathbf{X}^T \hat{\mathbf{W}} \mathbf{X} + \mathbf{I})^{-1} (\mathbf{X}^T \hat{\mathbf{W}} \mathbf{X} + d \mathbf{I}) \hat{\boldsymbol{\beta}}_{IGRM}, \quad (13)$$

where $0 < d < 1$. Regardless of d value, the MSE of the $\hat{\beta}_{IGLE}$ is smaller than that of $\hat{\beta}_{IGRM}$ because the MSE of $\hat{\beta}_{IGLE}$ is equal to [33]

$$\text{MSE}(\hat{\beta}_{IGLE}) = \tau \sum_{j=1}^p \frac{(\lambda_j + d)^2}{\lambda_j (\lambda_j + 1)^2} + (d - 1)^2 \sum_{j=1}^p \frac{\alpha_j^2}{(\lambda_j + 1)^2}. \quad (14)$$

4. Liu-type estimator

Alternative to Liu estimator, the Liu-type estimator was proposed by Liu [44] to overcome the problem of severe multicollinearity. The inverse Gaussian Liu-type estimator (IGLT) is defined as

$$\hat{\beta}_{IGLT} = (\mathbf{X}^T \hat{\mathbf{W}} \mathbf{X} + k \mathbf{I})^{-1} (\mathbf{X}^T \hat{\mathbf{W}} \mathbf{X} - d \mathbf{I}) \hat{\beta}_{IGRM}, \quad (15)$$

where $-\infty < d < \infty$ and $k \geq 0$. In Eq. (15), the parameter k can be used totally to control the conditioning of $\mathbf{X}^T \hat{\mathbf{W}} \mathbf{X} + k \mathbf{I}$. After the reduction of $\mathbf{X}^T \hat{\mathbf{W}} \mathbf{X} + k \mathbf{I}$ is reach a desirable level, then the expected bias that is generated can be corrected with the so-called bias correction parameter, d [45-49].

Liu [44] proved that, in terms of MSE, the Liu-type estimator has superior properties over ridge estimator. The MSE of $\hat{\beta}_{IGLT}$ is defined as

$$\text{MSE}(\hat{\beta}_{IGLT}) = \tau \sum_{j=1}^p \frac{(\lambda_j - d)^2}{\lambda_j (\lambda_j + k)^2} + (d + k)^2 \sum_{j=1}^p \frac{\alpha_j^2}{(\lambda_j + k)^2}. \quad (16)$$

5. Two-parameter estimator

Following Asar and Genç [50] and Huang and Yang [51] the two-parameter estimator in linear regression model is defined as:

$$\hat{\beta}_{TPE} = (\mathbf{X}^T \mathbf{X} + k \mathbf{I})^{-1} (\mathbf{X}^T \mathbf{X} + k d \mathbf{I}) \hat{\beta}_{OLS}, \quad (17)$$

where $0 < d < 1$ and $k \geq 0$. For IGRM, the two-parameter estimator (IGTP) is defined as:

$$\hat{\beta}_{IGTP} = (\mathbf{X}^T \hat{\mathbf{W}} \mathbf{X} + k \mathbf{I})^{-1} (\mathbf{X}^T \hat{\mathbf{W}} \mathbf{X} + k d \mathbf{I}) \hat{\beta}_{IGRM}. \quad (18)$$

It is obviously noted that the $\hat{\beta}_{IGTP}$ is a combination of two different estimators IGRR and IGLE. Furthermore, if $k = 1$, Eq. (18) will be the $\hat{\beta}_{IGLE}$ while if $k = 0$, Eq. (18) will be the $\hat{\beta}_{IGRM}$. Besides, when $d = 0$, then Eq. (18) will equal $\hat{\beta}_{IGRR}$.

In terms of MSE, the two-parameter estimator has superior properties over ML estimator. The MSE of $\hat{\beta}_{IGTP}$ is defined as

$$\text{MSE}(\hat{\beta}_{IGTP}) = \tau \sum_{j=1}^{p+1} \left[\frac{(\lambda_j + kd)^2}{\lambda_j (\lambda_j + k)^2} + k^2 (d - 1)^2 \frac{\alpha_j^2}{(\lambda_j + k)^2} \right]. \quad (19)$$

6. Real application

To demonstrate the usefulness of the shrinkage estimators in real application, we present here a chemistry dataset with $(n, p) = (65, 15)$, where n represents the number of imidazo[4,5-b] pyridine derivatives, which are used as anticancer compounds. While p denotes the number of molecular descriptors, which are treated as explanatory variables [52]. The response of interest is the biological activities (IC_{50}). Quantitative structure-activity relationship (QSAR) study has become a great deal of importance in chemometrics. The principle of QSAR is to model several biological activities over a collection of chemical compounds in terms of their structural properties [53]. Consequently, using of regression model is one of the most important tools for constructing the QSAR model.

First, to check whether the response variable belongs to the inverse Gaussian distribution, a Chi-square test is used. The result of the test equals to 5.2762 with p-value equals to 0.2601. It is indicated from this result that the inverse Gaussian distribution fits very well to this response variable. That is, the following model is set

$$\hat{y}_{IC_{50}} = \exp\left(\sum_{j=1}^{15} \mathbf{x}_j \hat{\beta}_j\right). \quad (20)$$

Second, to check whether there is a relationship among the explanatory variables or not, Figure 1 displays the correlation matrix among the 15 explanatory variables. It is obviously seen that there are correlations greater than 0.90 among MW, SpMaxA_D, and ATS8v ($r = 0.96$), between SpMax3_Bh(s) and ATS8v ($r = 0.92$), and between Mor21v with Mor21e ($r = 0.93$).

Third, to test the existence of multicollinearity after fitting the inverse Gaussian regression model using log link function and the estimated dispersion parameter is 0.00103, the eigenvalues of the matrix $\mathbf{X}^T \hat{\mathbf{W}} \mathbf{X}$ are obtained as 1.884×10^9 , 3.445×10^6 , 2.163×10^5 , 2.388×10^4 , 1.290×10^3 , 9.120×10^2 , 4.431×10^2 , 1.839×10^2 , 1.056×10^2 , 5525, 3231, 2631, 1654, 1008, and 1.115. The determined condition number $CN = \sqrt{\lambda_{\max} / \lambda_{\min}}$ of the data is 40383.035 indicating that the severe multicollinearity issue is exist.

The estimated inverse Gaussian regression coefficients and the estimated theoretical MSE values for the MLE, and the used estimators are listed in Table 1". According to Table 1, it is clearly seen that the IGTP has MSE values less than the MSE of the IGRM, in general. Moreover, the MSE of the IGTP estimator is the lowest among all estimators. Specifically, it can be seen that the MSE of IGTP estimator was about 44.24%, 39.17%, 32.62%, and 12.11% lower than that of IGRM, IGRR, IGLE, and IGLT, respectively.




Figure 1. Correlation matrix among the 15 explanatory variables of the real data.

Table 1: The estimated coefficients and MSE values of the used estimators

$\hat{\beta}$	Methods				
	IGRM	IGRR	IGLE	IGLT	IGTP
MW	1.002	0.744	0.835	0.731	0.841
IC3	1.237	0.977	1.087	0.969	2.005
SpMaxA_D	-1.102	-1.363	-1.269	-0.905	-1.304
ATS8v	-1.379	-1.67	-1.846	-1.126	-1.101
MATS7v	-1.219	-1.48	-1.386	-1.019	-1.421
MATS2s	-1.215	-1.476	-1.382	-1.015	-1.417
GATS4p	-1.237	-1.498	-2.405	-1.037	-1.439
SpMax8_Bh.p	2.506	2.145	2.309	2.707	2.304

SpMax3_	2.069	1.808	1.902	2.269	1.867
Bh.s.					
P_VSA_e	2.001	1.739	1.833	2.2	1.798
_3					
TDB08m	-2.103	-2.365	-2.27	-1.903	-2.305
RDF100m	1.571	1.309	1.403	1.77	1.368
Mor21v	-2.434	-2.695	-2.601	-2.235	-2.636
Mor21e	-2.352	-2.613	-2.519	-2.152	-2.554
HATS6v	2.211	1.95	2.044	2.411	2.009
MSE	3.295	2.258	1.823	1.658	1.215

7. Conclusions

In this paper, we presented a thorough review of literature regarding the biased estimators in inverse Gaussian regression model when the multicollinearity is existing. According to real data application, the two-parameter estimator has better performance than IGRM, IGRR, IGLE, and IGLT, in terms of MSE. In conclusion, the use of the two-parameter estimator is recommended when multicollinearity is present in the inverse Gaussians regression model.

References

1. Bhattacharyya, G.K. and A. Fries, Inverse Gaussian regression and accelerated life tests. Lecture Notes-Monograph Series, 1982. 2: p. 101-117.
2. Ducharme, G.R., Goodness-of-fit tests for the inverse Gaussian and related distributions. Test, 2001. 10(2): p. 271-290.
3. Folks, J.L. and A.S. Davis, Regression models for the inverse Gaussian distribution. Statistical Distributions in Scientific Work, 1981. 4(1): p. 91-97.
4. Fries, A. and G.K. Bhattacharyya, Optimal design for an inverse Gaussian regression model. Statistics & probability letters, 1986. 4(6): p. 291-294.
5. Heinzl, H. and M. Mittlböck, Adjusted R2 Measures for the Inverse Gaussian Regression Model. Computational Statistics, 2002. 17(4): p. 525-544.
6. Lemesko, B.Y., et al., Inverse Gaussian model and its applications in reliability and survival analysis, in Mathematical and Statistical Models and Methods in Reliability. 2010, Springer. p. 433-453.
7. Malehi, A.S., F. Pourmohammadi, and K.A. Angali, Statistical models for the analysis of skewed healthcare cost data: A simulation study. Health Economics Review, 2015. 5: p. 1-11.
8. Babu, G.J. and Y.P. Chaubey, Asymptotics and bootstrap for inverse Gaussian regression. Annals of the Institute of Statistical Mathematics, 1996. 48(1): p. 75-88.
9. Chaubey, Y.P., Estimation in inverse Gaussian regression: comparison of asymptotic and bootstrap distributions. Journal of Statistical Planning and Inference, 2002. 100(2): p. 135-143.
10. Wu, L. and H. Li, Variable selection for joint mean and dispersion models of the inverse Gaussian distribution. Metrika, 2011. 75(6): p. 795-808.
11. De Jong, P. and G.Z. Heller, Generalized linear models for insurance data. Vol. 10. 2008: Cambridge University Press Cambridge.
12. Wan, A.T.K., A note on almost unbiased generalized ridge regression estimator under asymmetric loss. Journal of Statistical Computation and Simulation, 2007. 62(4): p. 411-421.
13. Akdeniz, F. and S. Kaçiranlar, On the almost unbiased generalized Liu estimator and unbiased estimation of the bias and mse. Communications in Statistics - Theory and Methods, 1995. 24(7): p. 1789-1797.
14. El Karoui, N., On the impact of predictor geometry on the performance on high-dimensional ridge-regularized generalized robust regression estimators. Probability Theory and Related Fields, 2017. 170(1-2): p. 95-175.
15. Fallah, R., M. Arashi, and S.M.M. Tabatabaey, On the ridge regression estimator with sub-space restriction. Communications in Statistics - Theory and Methods, 2017. 46(23): p. 11854-11865.
16. Huang, H., J. Wu, and W. Yi, On the restricted almost unbiased two-parameter estimator in linear regression model. Communications in Statistics - Theory and Methods, 2016. 46(4): p. 1668-1678.
17. Kaçiranlar, S. and I. Dawoud, On the performance of the Poisson and the negative binomial ridge predictors. Communications in Statistics - Simulation and Computation, 2017: p. 0-0.
18. Li, Y. and H. Yang, On the Performance of the Jackknifed Modified Ridge Estimator in the Linear Regression Model with Correlated or Heteroscedastic Errors. Communications in Statistics - Theory and Methods, 2011. 40(15): p. 2695-2708.
19. Lopez Martin, M.D.M., et al., On the Selection of the Ridge and Raise Factors. Indian Journal of Science and Technology, 2017. 10(13): p. 1-8.

20. Månsson, K., On ridge estimators for the negative binomial regression model. *Economic Modelling*, 2012. 29(2): p. 178-184.
21. Månsson, K., B.M.G. Kibria, and G. Shukur, On Liu estimators for the logit regression model. *Economic Modelling*, 2012. 29(4): p. 1483-1488.
22. Månsson, K. and G. Shukur, On Ridge Parameters in Logistic Regression. *Communications in Statistics - Theory and Methods*, 2011. 40(18): p. 3366-3381.
23. Muniz, G. and B.M.G. Kibria, On Some Ridge Regression Estimators: An Empirical Comparisons. *Communications in Statistics - Simulation and Computation*, 2009. 38(3): p. 621-630.
24. Nomura, M., On exact small sample properties of the minimax generalized ridge regression estimators. *Communications in Statistics - Simulation and Computation*, 1988. 17(4): p. 1213-1229.
25. Ohtani, K., On small sample properties of the almost unbiased generalized ridge estimator. *Communications in Statistics - Theory and Methods*, 1986. 15(5): p. 1571-1578.
26. Segerstedt, B., On ordinary ridge regression in generalized linear models. *Communications in Statistics - Theory and Methods*, 1992. 21(8): p. 2227-2246.
27. Şiray, G.Ü., S. Toker, and S. Kaçiranlar, On the Restricted Liu Estimator in the Logistic Regression Model. *Communications in Statistics - Simulation and Computation*, 2014. 44(1): p. 217-232.
28. Toker, S. and S. Kaçiranlar, On the performance of two parameter ridge estimator under the mean square error criterion. *Applied Mathematics and Computation*, 2013. 219(9): p. 4718-4728.
29. van Wieringen, W.N., On the mean squared error of the ridge estimator of the covariance and precision matrix. *Statistics & Probability Letters*, 2017. 123: p. 88-92.
30. Varathan, N. and P. Wijekoon, On the Restricted Almost Unbiased Ridge Estimator in Logistic Regression. *Open Journal of Statistics*, 2016. 06(06): p. 1076-1084.
31. Wu, J., On the predictive performance of the almost unbiased Liu estimator. *Communications in Statistics - Theory and Methods*, 2015. 45(17): p. 5193-5203.
32. Wu, J. and H. Yang, On the Principal Component Liu-type Estimator in Linear Regression. *Communications in Statistics - Simulation and Computation*, 2014. 44(8): p. 2061-2072.
33. Kibria, B.M.G., K. Månsson, and G. Shukur, A Simulation Study of Some Biasing Parameters for the Ridge Type Estimation of Poisson Regression. *Communications in Statistics - Simulation and Computation*, 2015. 44(4): p. 943-957.
34. Baksalary, J.K., P.R. Pordzik, and G. Trenkler, A note on generalized ridge estimators. *Communications in Statistics - Theory and Methods*, 1990. 19(8): p. 2871-2877.
35. Firinguetti, L., H. Rubio, and Y.P. Chaubey, A non stochastic ridge regression estimator and comparison with the James-Stein estimator. *Communications in Statistics - Theory and Methods*, 2016. 45(8): p. 2298-2310.
36. Lazaridis, A., A Note Regarding the Condition Number: The Case of Spurious and Latent Multicollinearity. *Quality & Quantity*, 2007. 41(1): p. 123-135.
37. Özkale, M.R. and S. Kaçiranlar, A Prediction-Oriented Criterion for Choosing the Biasing Parameter in Liu Estimation. *Communications in Statistics - Theory and Methods*, 2007. 36(10): p. 1889-1903.
38. Shen, X., et al., A novel generalized ridge regression method for quantitative genetics. *Genetics*, 2013. 193(4): p. 1255-68.
39. Hoerl, A.E. and R.W. Kennard, Ridge regression: Biased estimation for nonorthogonal problems. *Technometrics*, 1970. 12(1): p. 55-67.
40. Asar, Y. and A. Genç, New shrinkage parameters for the Liu-type logistic estimators. *Communications in Statistics - Simulation and Computation*, 2015. 45(3): p. 1094-1103.
41. Algamal, Z.Y. and M.H. Lee, Penalized Poisson Regression Model using adaptive modified Elastic Net Penalty. *Electronic Journal of Applied Statistical Analysis*, 2015. 8(2): p. 236-245.
42. Uusipaikka, E., Confidence intervals in generalized regression models. 2009, NW: Chapman & Hall/CRC Press.
43. Månsson, K. and G. Shukur, A Poisson ridge regression estimator. *Economic Modelling*, 2011. 28(4): p. 1475-1481.
44. Liu, K., Using Liu-type estimator to combat collinearity. *Communications in Statistics-Theory and Methods*, 2003. 32(5): p. 1009-1020.
45. Alheety, M.I. and B.M. Golam Kibria, Modified Liu-Type Estimator Based on $(r - k)$ Class Estimator. *Communications in Statistics - Theory and Methods*, 2013. 42(2): p. 304-319.
46. Kibria, B.M.G. and A.K.M.E. Saleh, Preliminary test ridge regression estimators with student's t errors and conflicting test-statistics. *Metrika*, 2004. 59(2): p. 105-124.
47. Norouzrad, M. and M. Arashi, Preliminary test and Stein-type shrinkage ridge estimators in robust regression. *Statistical Papers*, 2017.

48. Wu, J., Preliminary test Liu-type estimators based on W, LR, and LM test statistics in a regression model. Communications in Statistics - Simulation and Computation, 2016. 46(9): p. 6760-6771.
49. Wu, J., Modified Restricted Almost Unbiased Liu Estimator in Linear Regression Model. Communications in Statistics - Simulation and Computation, 2014. 45(2): p. 689-700.
50. Asar, Y. and A. Genç, A New Two-Parameter Estimator for the Poisson Regression Model. Iranian Journal of Science and Technology, Transactions A: Science, 2017.
51. Huang, J. and H. Yang, A two-parameter estimator in the negative binomial regression model. Journal of Statistical Computation and Simulation, 2014. 84(1): p. 124-134.
52. Algamal, Z.Y., et al., High-dimensional QSAR prediction of anticancer potency of imidazo[4,5-b]pyridine derivatives using adjusted adaptive LASSO. Journal of Chemometrics, 2015. 29(10): p. 547-556.
53. Algamal, Z.Y. and M.H. Lee, A novel molecular descriptor selection method in QSAR classification model based on weighted penalized logistic regression. Journal of Chemometrics, 2017. 31(10): p. e2915.

مقدرات التقليص في نموذج الانحدار الكاوسي المعكوس: مراجعة الموضوع

فرح عبد الغني يونس و رفل اديب عثمان

قسم الاحصاء والمعلوماتية، كلية علوم الحاسوب والرياضيات، جامعة الموصل، الموصل، العراق.

الخلاصة : ان وجود علاقة ارتباط عالية بين المتنبئين في نمذجة الانحدار له تأثيرات غير مرغوب فيها على تقدير الانحدار. هناك العديد من الطرق المتحيزة المتاحة للتغلب على هذه المشكلة. نموذج الانحدار الغاوسي العكسي (IGRM) هو نموذج خاص من النماذج الخطية المعممة. يعد IGRM نموذجًا معروفًا في تطبيق البحث عندما يكون متغير الاستجابة تحت الدراسة عبارة عن بيانات منحرفة. تم اقتراح العديد من المقدرات المتحيزة للتغلب على العلاقة الخطية المتعددة في IGRM في الأدبيات باستخدام نظريات مختلفة. يتم تقديم لمحة عامة عن الأساليب المتحيزة الحديثة لـ IGRM. تسمح لنا المقارنة بين هذه المقدرات المتحيزة بالحصول على نظرة ثاقبة لأدائها. **الكلمات المفتاحية :** متعدد الخطية. مقرر متحيز نموذج الانحدار الغاوسي المعكوس ؛ محاكاة مونت كارلو